

# BOYLE'S LAW

Name \_\_\_\_\_

Boyle's Law states that the volume of a gas varies inversely with its pressure if temperature is held constant. (If one goes up, the other goes down.) We use the formula:

$$P_1 \times V_1 = P_2 \times V_2$$

Solve the following problems (assuming constant temperature).

1. A sample of oxygen gas occupies a volume of 250. mL at 740. torr pressure. What volume will it occupy at 800. torr pressure?  
\_\_\_\_\_
2. A sample of carbon dioxide occupies a volume of 3.50 liters at 125 kPa pressure. What pressure would the gas exert if the volume was decreased to 2.00 liters?  
\_\_\_\_\_
3. A 2.0 liter container of nitrogen had a pressure of 3.2 atm. What volume would be necessary to decrease the pressure to 1.0 atm?  
\_\_\_\_\_
4. Ammonia gas occupies a volume of 450. mL at a pressure of 720. mm Hg. What volume will it occupy at standard pressure?  
\_\_\_\_\_
5. A 175 mL sample of neon had its pressure changed from 75 kPa to 150 kPa. What is its new volume?  
\_\_\_\_\_
6. A sample of hydrogen at 1.5 atm had its pressure decreased to 0.50 atm producing a new volume of 750 mL. What was its original volume?  
\_\_\_\_\_
7. Chlorine gas occupies a volume of 1.2 liters at 720 torr pressure. What volume will it occupy at 1 atm pressure?  
\_\_\_\_\_
8. Fluorine gas exerts a pressure of 900. torr. When the pressure is changed to 1.50 atm, its volume is 250. mL. What was the original volume?  
\_\_\_\_\_

# CHARLES' LAW

Name \_\_\_\_\_

Charles' Law states that the volume of a gas varies directly with the Kelvin temperature assuming that pressure is constant. We use the following formulas:

$$\frac{V_1}{T_1} = \frac{V_2}{T_2} \quad \text{or} \quad V_1 \times T_2 = V_2 \times T_1$$
$$K = ^\circ C + 273$$

Solve the following problems assuming a constant pressure.

1. A sample of nitrogen occupies a volume of 250 mL at 25° C. What volume will it occupy at 95° C?  
\_\_\_\_\_
2. Oxygen gas is at a temperature of 40° C when it occupies a volume of 2.3 liters. To what temperature should it be raised to occupy a volume of 6.5 liters?  
\_\_\_\_\_
3. Hydrogen gas was cooled from 150° C to 50° C. Its new volume is 75 mL. What was its original volume?  
\_\_\_\_\_
4. Chlorine gas occupies a volume of 25 mL at 300 K. What volume will it occupy at 600 K?  
\_\_\_\_\_
5. A sample of neon gas at 50° C and a volume of 2.5 liters is cooled to 25° C. What is the new volume?  
\_\_\_\_\_
6. Fluorine gas at 300 K occupies a volume of 500 mL. To what temperature should it be lowered to bring the volume to 300 mL?  
\_\_\_\_\_
7. Helium occupies a volume of 3.8 liters at -45° C. What volume will it occupy at 45° C?  
\_\_\_\_\_
8. A sample of argon gas is cooled and its volume went from 380 mL to 250 mL. If its final temperature was -55° C, what was its original temperature?  
\_\_\_\_\_